





POSITIONING STATEMENT

ageLOC TR90 JumpStart is designed to be used as part of the TR90 program, a weight management and body shaping system designed to promote a leaner, younger looking you.* JumpStart contains the powerful ageLOC JumpStart proprietary blend to support a healthy start to your weight management goals.

CONCEPT

When beginning a weight management system, a good start is key to your success. ageLOC TR90 JumpStart is the perfect push to get you primed and ready for success. You can see the beginnings of success in just 15 days—and experience all the motivation that comes along with it.* For the first 15 days of the ageLOC TR90 program, take this innovative product in the morning to help you achieve your goals. Mix it in water or your favorite beverage (for example, try it in hot water or q3).

PRIMARY BENEFITS

- Positively impacts gene expression*
- Formulated to get you off to a positive start

FREQUENTLY ASKED QUESTIONS

What makes ageLOC TR90 JumpStart unique? ageLOC TR90 JumpStart is a proprietary blend specially formulated with premium plant-based nutrients including prickly pear, saffron, pomegranate, and red orange.

Do I need to take ageLOC TR90 JumpStart at a specific time of day or in conjunction with a meal? ageLOC TR90 JumpStart should be taken each morning during the first 15 days of the TR90 program. It does not need to be taken in conjunction with a meal. If you forget to

take it in the morning, or if you simply prefer to take it at a different time of the day, it is suitable to take at any time of the day.

Can I take ageLOC TR90 JumpStart on its own? ageLOC TR90 JumpStart can be taken on its own, but it's recommended to be used with the entire TR90 system for best results.

Can ageLOC TR90 Jumpstart be taken longer than 15 days and is it safe?

ageLOC TR90 Jumpstart was specifically formulated to provide an initial boost to the benefits of the program, but can also be used safely for the duration of the full TR90 program for individuals seeking additional support.

KEY SCIENTIFIC STUDIES

- Titta L, Trinei M, Stendardo M, Berniakovich I, Petroni K, Tonelli C, Riso P, Porrini M, Minucci S, Pelicci PG, Rapisarda P, Reforgiato Recupero G, Giorgio M. Blood orange juice inhibits fat accumulation in mice. Int J Obes (Lond). 2010 Mar;34(3):578-88. doi: 10.1038/ijo.2009.266. Epub 2009 Dec 22. PubMed PMID: 20029381.
- 2. Gout B, Bourges C, Paineau-Dubreuil S. Satiereal, a Crocus sativus L extract, reduces snacking and increases satiety in a randomized placebo-controlled study of mildly overweight, healthy women. Nutr Res. 2010 May;30(5):305-13. doi: 10.1016/j.nutres.2010.04.008. PubMed PMID: 20579522.
- 3. González-Ortiz M, Martínez-Abundis E, Espinel-Bermúdez MC, Pérez-Rubio KG. Effect of pomegranate juice on insulin secretion and sensitivity in patients with obesity. Ann Nutr Metab. 2011;58(3):220-3. doi: 10.1159/000330116. Epub 2011 Jul 28. PubMed PMID: 21811060.

(C) ageloc TR90 JUMPSTART

- Tesoriere L, Butera D, Pintaudi AM, Allegra M, Livrea MA. Supplementation with cactus pear (Opuntia ficus-indica) fruit decreases oxidative stress in healthy humans: a comparative study with vitamin C. Am J Clin Nutr. 2004 Aug;80(2):391-5. PubMed PMID: 15277160.
- Godard MP, Ewing BA, Pischel I, Ziegler A, Benedek B, Feistel B. Acute blood glucose lowering effects and long-term safety of OpunDia supplementation in prediabetic males and females. J Ethnopharmacol. 2010 Aug 9;130(3):631-4. doi: 10.1016/j.jep.2010.05.047. Epub 2010 Jun 4. PubMed PMID: 20621660.

AGELOC TR90 JUMPSTART

cts	
Amount % Daily Per Serving Value*	
5 2 g 1%** < 1 g 3%**	
2000 mg *	
177 mg *	
150 mg *	
125 mg *	
	Per Serving Value* 5 2 g 1%** <1 g 3%** 2000 mg * 177 mg * 150 mg *

USAGE

Mix one (1) packet in two to eight (2-8) ounces of water or favorite beverage. Drink each morning for the first 15 days. Store in a cool, dry place.

WARNINGS

Keep out of reach of children. Consult a physician prior to use if pregnant or lactating, or prior to using prescription medication. Discontinue use 2 weeks prior to and after surgery. Discontinue use and consult a physician if any adverse reactions occur. Do not use if safety seal is broken or missing.

*This statement has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

